




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Asthma in older adults with severe COVID-19: clinical outcomes and predictors of mortality.

Sergey N. Avdeev MD, PhD , Anna A. Pozdnakova MD ,
Viliya V. Gaynitdinova MD, PhD , Svetlana Yu Chikina MD, PhD ,
Anna E. Vlasenko PhD

PII: S1081-1206(21)01163-7
DOI: <https://doi.org/10.1016/j.anai.2021.10.016>
Reference: ANAI 3744



To appear in: *Annals of Allergy, Asthma Immunology*

Received date: 8 August 2021
Revised date: 7 October 2021
Accepted date: 11 October 2021

Please cite this article as: Sergey N. Avdeev MD, PhD , Anna A. Pozdnakova MD , Viliya V. Gaynitdinova MD, PhD , Svetlana Yu Chikina MD, PhD , Anna E. Vlasenko PhD , Asthma in older adults with severe COVID-19: clinical outcomes and predictors of mortality., *Annals of Allergy, Asthma Immunology* (2021), doi: <https://doi.org/10.1016/j.anai.2021.10.016>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Asthma in older adults with severe COVID-19: clinical outcomes and predictors of mortality.

Sergey N. Avdeev¹, MD, PhD

Anna A. Pozdnakova², MD

Viliya V. Gaynitdinova¹, MD, PhD

Svetlana Yu. Chikina¹, MD, PhD

Anna E. Vlasenko³, PhD

¹Department of Pulmonology, I.M. Sechenov First Moscow State Medical University (Sechenov University), Moscow, Russia.

²University Clinical Hospital №4, I.M. Sechenov First Moscow State Medical University (Sechenov University), Moscow, Russia.

³Department of Medical Cybernetics and Informatics, Novokuznetsk State Institute for Advanced Training of Doctors, Novokuznetsk, Russia.

Corresponding author:

Prof. Dr. Sergey N. Avdeev

Department of Pulmonology

I.M. Sechenov First Moscow State Medical University (Sechenov University)

Ministry of Health of the Russian Federation

8, Trubetskaya Street, Moscow, 119991, Russia

Tel: +7(495)708-35-76

E-mail: serg_avdeev@list.ru

<http://orcid.org/0000-0002-5999-2150>

Conflicts of interest:

The authors declare that they have no relevant conflicts of interest.

Funding Source:

None.

Keywords:

asthma, elderly, COVID-19, SARS-CoV-2

Abbreviations:

CCI, Charlson comorbidity index

COVID-19, Coronavirus disease

ICU, intensive care unit

SpO₂, oxygen saturation

Total word count:

990.

Figures:

0

Tables:

1.

Contributorship Statement: All authors were involved in the assessment and treatment of the patient, and in the writing and correction of the manuscript.

Ethical approval

The study was approved by the Medical Ethical Committee of Sechenov Moscow Medical University (protocol number 08-20/5).

To date, numerous studies have examined the relationship between COVID-19 and asthma; however, the impact of asthma on COVID-19-associated outcomes remains controversial and not fully understood¹⁻³. It is possible that these contradictions can be explained in terms of asthma heterogeneity. There is data to suggest that asthma in older adults is phenotypically different from that in young patients, particularly when physiological changes associated with the aging process are taken into account⁴. Since old age is a well-known factor of a poor prognosis in COVID-19⁵, it is of great interest to study the course of COVID-19 in older patients with asthma.

This observational study was conducted in the Pulmonology Department of a university-affiliated hospital between May 12, 2020, and December 27, 2020. As this was a retrospective study, the requirement for informed consent was waived. We enrolled patients over 65 years old with physician diagnosed asthma admitted with SARS-CoV-2 infection confirmed by real-time polymerase chain reaction, radiological findings compatible with severe COVID-19 pneumonia, and SpO₂ at admission <92% (on room air). The exclusion criteria were non-asthma chronic lung diseases. The diagnostic criteria of allergic asthma were those of the GINA guidelines: an exposure to the allergen induces or aggravates the symptoms, and the skin prick test and serum specific IgE test show positive responses to at least one allergen⁶. Demographic, clinical, and laboratory data was recorded at admission. We recorded the Charlson Comorbidity Index (CCI), which calculates the score for comorbidities according to the relative risks of 19 major diseases, including ischemic heart disease, diabetes, and hypertension. We also analyzed the outcomes of COVID-19, such as a transfer to the intensive care unit (ICU), need for non-invasive and invasive mechanical ventilation, and 28-day mortality.

The objectives of this study were: (1) to investigate the influence of asthma in older adults on COVID-19 outcomes, and (2) to identify predictors of in-hospital mortality of older patients

with asthma and COVID-19. The Cox proportional hazard models were used to estimate the association between asthma and in-hospital mortality; the models were adjusted for age, sex, and comorbidities.

Of the 2,435 patients hospitalized with COVID-19 infection, 69 patients met the necessary diagnostic criteria (2.8%). Baseline characteristics of the study population are shown in Table 1. Most patients had a non-allergic asthma phenotype (44 patients, 63.8%) and multiple comorbidities (the median CCI was 5.3). Nineteen of included patients (27.5%) were receiving GINA step 4-5 therapies; 12 patients (17.4%) and 4 patients (5.8%) were treated before admission with maintenance oral steroids and biologic therapy ((omalizumab (n=2), mepolizumab (n=2)), respectively.

During hospitalization, all patients received the standard therapy including prophylactic enoxaparin, intravenous dexamethasone, and tocilizumab. Supplemental oxygen was administered in all patients (100%), non-invasive ventilation and invasive mechanical ventilation were used in 6 (8.6%) and 7(10.1%) patients, respectively. Twelve patients (17.3%) were transferred to the ICU, and the 28-day mortality rate was 13%(9 patients). All patients who died had non-allergic asthma. There was a significant difference in the proportion of patients with severe asthma among deceased and surviving patients (56% vs 13%; $p=0.01$). Patients who died from COVID-19 used maintenance oral steroids more often than survivors (44% vs 13%, $p=0.04$).

Patients who died had a significantly higher CCI, higher body temperature at admission, higher respiratory rate, and a higher baseline blood neutrophil count, neutrophil/lymphocyte ratio, fibrinogen, lactate dehydrogenase, glucose levels, and C-reactive protein at Day 5 compared to survivors. The Cox regression analysis identified the following variables predicting poor outcomes: CCI (risk ratio (RR) 1.67[1.11-2.53]), body temperature (RR 3.73[1.31-10.6]), severe asthma (RR 2.75[1.08-6.99]), non-allergic asthma (RR 7.95[1.06-

14.30]), and the chronic oral steroid use (RR 2.57[1.01-6.53]). The risk of death was not associated with an increased or decreased eosinophil count (Cox regression; $p=0.57$ and $p=0.53$ for percentage and absolute values, respectively).

In a recently published study, Eggert and colleagues showed that among asthmatics positive for SARS-CoV-2, the allergic asthma phenotype was associated with a reduced risk of hospital admissions, meaning that certain phenotypes of asthma may be protective⁷. The role of allergic asthma diminishes with age; hence a different phenotype presents in older adults, with normal eosinophils and elevated sputum neutrophils⁴. Interestingly, in our study, all deceased patients had a non-allergic asthma phenotype. In addition, as recently shown in a study by Schleich et al., older adults with asthma are characterized by poorer lung function and higher bronchial neutrophilic inflammation⁴. Recent publications have suggested that allergic status and type 2 inflammation may decrease susceptibility to SARS-CoV-2 infection and protect against the most severe consequences of COVID-19⁸. Ferastraoaru et al. in their retrospective study demonstrated that in patients with asthma pre-existing blood eosinophilia greater than or equal to 150 cells/mL was protective from COVID-19-associated admission⁹. In our cohort of older adults with asthma, we were unable to show differences in eosinophilia levels between deceased and surviving patients. However, it cannot be ruled out that the blood eosinophil counts did not differ significantly due to the outpatient and inpatient use of systemic steroids. In our study therapy with maintenance oral steroids in older adults with asthma was a risk factor for poor prognosis. Similar data was presented in a recent study by Adir et al. who also showed that systemic steroid use was significantly associated with an increased risk of severe COVID-19 and all-cause mortality¹⁰. In contrast to the study by Adir et al., in our study, all patients had severe COVID-19 and all received intravenous dexamethasone during hospitalization. Despite this, a chronic use of steroids turned out to be a factor of poor prognosis.

Our study has several limitations. It was a retrospective study performed in a single center, and statistical analysis and interpretation of our study results are further limited by its relatively small sample size.

Therefore, our study provided evidence that non-allergic asthma phenotype, asthma severity, systemic steroid use, and comorbidities may be risk factors for poor outcomes in older patients with asthma and severe COVID-19. However, further large cohort studies are needed to examine the effect of asthma severity and phenotypes on COVID-19 outcomes in older adults.

References

1. Heffler E, Detoraki A, Contoli M, et al. COVID-19 in severe asthma network in Italy (SANI) patients: clinical features, impact of comorbidities and treatments. *Allergy*. 2021;76(3):887–892.
2. Rosenthal JA, Awan SF, Fintzi J, et al. Asthma is associated with increased risk of intubation but not hospitalization or death in coronavirus disease 2019. *Ann Allergy Asthma Immunol*. 2021;126(1):93–95.
3. Avdeev S, Moiseev S, Brovko M, et al. Low prevalence of bronchial asthma and chronic obstructive lung disease among intensive care unit patients with COVID-19. *Allergy*. 2020;75(10):2703–2704.
4. Schleich F, Graff S, Guissard F, et al. Asthma in elderly is characterized by increased sputum neutrophils, lower airway caliber variability and air trapping. *Respir Res*. 2021;22(1):15.
5. Cummings MJ, Baldwin MR, Abrams D, et al. Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet*. 2020;395(10239):1763–1770.

6. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention, 2021. Available from: www.ginasthma.
7. Eggert LE, He Z, Collins W, et al. Asthma phenotypes, associated comorbidities, and long-term symptoms in COVID-19. *Allergy*. 2021 Jun 3;10.1111/all.14972. doi: 10.1111/all.14972. Epub ahead of print.
8. Jackson DJ, Busse WW, Bacharier LB, et al. Association of respiratory allergy, asthma, and expression of the SARS-CoV-2 receptor ACE2. *J Allergy Clin Immunol*. 2020;146(1):203–206.e3.
9. Ferastraoaru D, Hudes G, Jerschow E, et al. Eosinophilia in asthma patients is protective against severe COVID-19 illness. *J Allergy Clin Immunol Pract*. 2021;9(3):1152–1162.e3.
10. Adir Y, Humbert M, Saliba W. COVID-19 risk and outcomes in adult asthmatic patients treated with biologics or systemic corticosteroids: Nationwide real-world evidence. *J Allergy Clin Immunol*. 2021;148(2):361–367.e13.

Table 1. Demographic, clinical and laboratory characteristics of older patients with asthma and severe COVID-19.

Parameters	All patients (n=69)	Survived patients (n=60)	Deceased patients (n=9)	P value ¹
Age, years	71(67-80)	70(66-80)	72(68-85)	0.33
Gender (male/female)	30/39	24/36	6/3	0.16
Smoking history, pack-years	45(32-50)	48(45-55)	32(20-45)	0.52
BMI, kg/m ²	28.2(24.7-33.2)	28.1(23.2-33.8)	28.0(27.3-32.4)	0.69
CCI, points	5.3(4-7)	5.0(3-6)	7.0(6-7.5)	0.003
Arterial hypertension, n(%)	31(44.9%)	25(41.7%)	6(66.7%)	0.30
Ischemic heart disease, n(%)	18(26.1%)	14(23.3%)	4(44.4)	0.26
Atrial fibrillation, n(%)	25(36.2%)	20(29.0%)	5(55.6%)	0.52
Obesity, n(%)	30(43.5%)	25(41.7%)	5(55.6%)	0.42
Diabetes mellitus, n(%)	20(28.9%)	17(28.3%)	3(33.3%)	0.54

Chronic kidney disease, n(%)	9(13.0%)	6(10.0%)	3(33.3%)	0.14
Allergic asthma phenotype, n(%)	22(31.9%)	22(36.7%)	0(0%)	0.05
Oral steroids, n(%)	12(17.4%)	8(13.3%)	4(44.4%)	0.04
Biologic therapy, n(%)	4(5.8%)	4(6.7%)	0(0%)	0.58
Body temperature, °C*	37.1(36.9-37.6)	37.0(36.8-37.5)	37.8(37.5-38.5)	0.006
Respiratory rate, min ⁻¹	23(22-24)	23(22-24)	25(24-26)	0.02
Heart rate, min ⁻¹	83(74-94)	82(74-95)	84(75-90)	0.57
SpO ₂ /FiO ₂	207(175-229)	209(171-229)	206(187-323)	0.64
Leukocytes, 10 ⁹ /L	6.2(4.7-8.9)	6.1(4.6-8.2)	8.9(6.1-9.3)	0.13
Neutrophils, 10 ⁹ /L	4.6 (2.8-10.8)	3.8(2.8-6.2)	7.3(5.1-8.2)	0.05
Lymphocytes, 10 ⁹ /L	0.9 (0.6-1.4)	1.0(0.7-1.4)	0.7(0.4-0.9)	0.06
Neutrophil/lymphocyte ratio	4.5(2.6-10.3)	3.5(2.3-7.2)	10.7(7.1-11.2)	0.02
Eosinophils, %	0.6(0.3-2.0)	1.0(0.3-3.0)	0.4(0.2-1.3)	0.15
Eosinophils, 10 ⁹ /L	0.1(0.07-0.2)	0.1(0.03-0.2)	0.1(0.07-0.1)	0.46
Fibrinogen, g/L	5.1(4.2-8.9)	4.9(3.9-7.8)	11.4(10.3-12.4)	0.04
D-dimer, mg/L	0.9(0.4-1.1)	0.7(0.4-1.2)	1.0(0.8-5.0)	0.21
CRP at admission, mg/L	40.0(14.4-73.2)	34.9(10.6-70.9)	50.0(35.1-105.9)	0.17
CRP at day 5, mg/L	17.7(3.5-30.2)	8.6(2.8-24.3)	39.1(20.3-170.9)	0.007
Creatinine, µmol/L	90.8(80.2-103.3)	90.2(78.4-100.1)	97.8(82.4-110.9)	0.39
Glucosa, mmol/L	6.6(5.7-8.1)	6.5(5.5-7.8)	8.0(7.3-10.2)	0.02
ALT, U/L	28.7(17.7-42.8)	30.4(16-44)	24.5(19.2-58)	0.79
AST, U/L	34.4(25-49.6)	31.5(24.9-47.3)	36.5(28-60.5)	0.59
LDH, U/L	536(399-694)	445(382-664)	701(532-1155)	0.03
CT, % of lung involvement	30(25-70)	30(25-50)	50(25-75)	0.66

*The highest temperature during the day was recorded.

Continuous variables are presented as median value [interquartile range (IQR)].

Abbreviations: BMI, body mass index; CCI, Charlson comorbidity index; SpO₂, oxygen saturation; FiO₂, inspired oxygen fraction; C-reactive protein; ALT, alanine aminotransferase; AST, aspartate aminotransferase; LDH, lactate dehydrogenase; CT, computed tomography.

¹Mann-Whitney test was used.